

**Effects of Epstein-Barr Virus on inflammatory cytokines production in gingival
fibroblasts and RANKL-induced osteoclast differentiation in RAW264.7 cells**

Sho Yokoe

Nihon University Graduate School of Dentistry,

Major in Periodontology

(Directors: Prof. Shuichi Sato and Prof. Kenichi Imai)

Contents

	Page
Abstract1
Introduction2
Materials and Methods5
Results10
Discussion13
Conclusion17
References18
Figures22

This doctoral thesis was prepared using the original article “Epstein-Barr Virus promotes the production of inflammatory cytokines in gingival fibroblasts and RANKL-induced osteoclast differentiation in RAW264.7 cells” (Sho Yokoe, Akira Hasuike, Norihisa Watanabe, Hideki Tanaka, Yukihiro Karahashi, Shin Wakuda, Osamu Takeichi, Takayuki Kawato, Hideki Takai, Yorimasa Ogata, Shuichi Sato, Kenichi Imai. *Int. J. Mol. Sci.* 2022, 23, 809.) with new unpublished data (Fig. 4C and 4D).

Abstract

Periodontitis is an inflammatory condition that causes the destruction of the supporting tissues of teeth and is a major public health problem affecting more than half of the adult population worldwide. Recently, members of the herpes virus family, such as Epstein–Barr virus (EBV), have been suggested to be involved in the etiology of periodontitis because bacterial activity alone does not adequately explain the clinical characteristics of periodontitis.

Many studies have found a higher prevalence of EBV DNA in the gingival tissue, gingival crevicular fluid, and subgingival plaque of patients with periodontitis. Additionally, previous work also reported that EBV DNA was more frequently detected in patients with deeper probing depth sites than in those with shallow probing depth sites or healthy controls.

However, the role of EBV in the etiology of periodontitis is unknown. This study aimed to examine the effect of inactivated EBV on the expression of inflammatory cytokines in human gingival fibroblasts (HGFs) and induction of osteoclast differentiation. The study found that extremely high levels of interleukin (IL)-6 and IL-8 were induced by inactivated EBV in a copy-dependent manner in HGFs. The levels of IL-6 and IL-8 in HGFs were higher when the cells were treated with EBV than when treated with lipopolysaccharide and lipoteichoic acid. EBV induced I κ B α degradation, NF- κ B transcription, and RAW264.7 cell differentiation into osteoclast-like cells.

These findings suggest that even without infecting the cells, EBV contributes to inflammatory cytokine production and osteoclast differentiation by contact with oral cells or macrophage lineage, resulting periodontitis onset and progression.

Introduction

Periodontitis is a chronic inflammatory disease that causes the destruction of periodontal tissues and alveolar bone and is among the most prevalent infectious diseases in the world. Although no single etiological agent has been identified, a number of bacteria, such as *Porphyromonas gingivalis*, are believed to be associated with the disease and are used as diagnostic markers (1). However, bacterial activity alone cannot explain the several clinical characteristics of periodontitis (2). Several reports have demonstrated the absence of periodontal bacteria in patients with periodontitis, and no significant difference has been found in the prevalence of bacteria between healthy and diseased periodontium (3–5).

Recently, a positive association has been reported between periodontitis and infection with herpesviruses, such as Epstein–Barr virus (EBV) and human cytomegalovirus (CMV) (2, 6). EBV is an enveloped herpes virus with double-stranded DNA and only infects humans. During a primary EBV infection, the virus undergoes lytic replication in epithelial cells and B cells, in which it establishes latency (7). Periodically, the virus can reactivate, and it is commonly found in the saliva of infected persons (2).

Recently, previous work demonstrated that *P. gingivalis* can induce EBV reactivation through epigenetic regulation (8). Many studies have found a higher prevalence of EBV DNA in the gingival tissue, gingival crevicular fluid, and subgingival plaque of patients with periodontitis (9–13). Additionally, previous work also reported that EBV DNA was more frequently detected in patients with deeper probing depth sites than in those with shallow probing depth sites or healthy controls. The range of counts of EBV DNA in the periodontal pockets of periodontitis patients was 3.74×10^3 – 2.83×10^9 copies/mL (14, 15). A large number of EBV-encoded small RNA (EBER) positive cells are found in the gingival tissue of

patients with periodontal disease (14). Vincent-Bugnas et al. reported that the level of gingival epithelial EBV infection is correlated with the severity of periodontitis (16).

Treatment with an antiherpesvirus drug decreased the presence of EBV to the detection limit, and the periodontal condition improved dramatically (17). These observations indicate that EBV is involved in the etiology of periodontal disease. However, despite its importance, the molecular mechanisms by which EBV affects the pathogenesis and progression of periodontal disease remain poorly understood.

It has been demonstrated that periodontopathic bacterial infections and their associated bacterial products, such as lipopolysaccharide (LPS), stimulate the production of inflammatory cytokines such as IL-6 and IL-8 from human gingival fibroblasts (HGFs) and monocytes, leading to connective tissue destruction and bone resorption in periodontal patients (1, 18). Both LPS and inflammatory cytokines promote osteoclastogenesis and enhance osteoclast-mediated bone resorption by promoting the expression of the receptor activator of nuclear factor kappa B (NF- κ B) ligand (RANKL) (1, 18, 19).

EBV can also trigger the cells of the immune system to produce cytokines and monokines, and inflammatory mediators play an important role in the pathogenesis and eventual outcome of EBV infection (7). Although the main cells targeted by EBV infection are B cells, several studies have reported that UV- or heat-inactivated EBV could stimulate the production of inflammatory cytokines in monocytes and neutrophils (20–23). These observations indicate that that even when EBV has not infected the cells, simple contact with host cells can induce the production of inflammatory cytokines that are involved in the progression of EBV infection.

The purpose of this study is to examine whether inactivated EBV affected the expression of inflammatory cytokines in HGFs. HGFs are the major cells in periodontal connective tissues, providing a tissue framework for tooth anchorage and functioning as a regulator of

the production of inflammatory molecules. This study also investigated the potential of inactivated EBV for inducing osteoclast differentiation.

To the best of recognized knowledge, this is the first time that inactivated EBV has been shown to markedly induce the production of inflammatory cytokines and the activation of nuclear factor kappa B (NF- κ B) in oral cells and the differentiation of RAW264.7 cells into osteoclast-like cells.

Materials and Methods

Cell culture

HGFs were maintained at 37°C in Dulbecco's modified Eagle's medium (Sigma Aldrich, St. Louis, MO, USA) with 10% heat-inactivated fetal bovine serum (FBS) (Thermo Fisher Scientific, Rockford, IL, USA), penicillin (100 U/mL), and streptomycin (100 µg/mL). The Ethics Committee of Nihon University School of Dentistry at Matsudo approved the study (EC03-041, EC10-040, and EC14-023), and written informed consent was obtained from each patient after full explanation of the protocol. The mouse macrophage-like cell line RAW264.7 was purchased from Dainippon Pharmaceutical (Osaka, Japan) and maintained in α -minimal essential medium (WAKO, Osaka, Japan) supplemented with 10% FBS and penicillin/streptomycin.

Reagents and plasmids

Inactivated EBV (strain: B95-8) and CMV (strain: AD-169) were purchased from ZeptoMetrix (Buffalo, NY, USA). These viruses were isolated from cell culture medium or patient plasma and were purified to a high degree. The viral surface proteins were chemically or enzymatically processed while leaving nucleic acids intact. A virus-treated solution (ZeptoMetrix, Buffalo, NY, USA) was added to the cells as a control. Human recombinant tumor necrosis factor- α (TNF- α) was purchased from Roche Diagnostics Deutschland GmbH (Mannheim, Germany). *P. gingivalis* LPS was obtained from Invivogen (St. Louis, MO, USA). *Escherichia coli* LPS and *Staphylococcus aureus* lipoteichoic acid (LTA) were purchased from Sigma Aldrich. Antibodies for, I κ B α , p65, PCNA and GAPDH were purchased from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA, USA). BAY11-7082, an inhibitor of I κ B- α phosphorylation, was purchased from Sigma Aldrich. The reporter plasmid

used, which expresses firefly luciferase under the control of NF- κ B (pGL3- κ B luc), was described previously (24, 25).

Preparation of mRNA and real-time polymerase chain reaction

The experimental procedures for RNA purification and real-time polymerase chain reaction (RT-PCR) were performed as previously described (25). Briefly, HGFs were washed once with ice-cold phosphate-buffered saline (PBS) and homogenized with QIAshredder (Qiagen, Hilden, Germany), and total RNA was purified using RNeasy Mini Kits (Qiagen). For cDNA synthesis, 1 μ g of total RNA was reverse transcribed using RNA PCR kits (PrimeScript; Takara Bio, Shiga, Japan). The resulting cDNA mixture was subjected to RT-PCR analysis using SYBR Premix Ex Taq solution (Takara Bio) containing 10 μ M sense and antisense primers. The primer sequences for each amplified gene were as follows: IL-6, forward (5'-CCA TAC CAG GTG CCT TTT GT-3') and reverse (5'-GAG ACT GGG AAC AGC TGA GG-3'); IL-8, forward (5'-CCA TAC CAG GTG CCT TTT GT-3') and reverse (5'-GAG ACT GGG AAC AGC TGA GG-3'); and GAPDH, forward (5'-ACC AGC CCC AGC AAG AGC ACA AG-3') and reverse (5'-TTC AAG GGG TCT ACA TGG CAA CTG-3'). The PCR assays were performed using TP-800 Thermal Cycler Dice Real-Time System (Takara Bio) and were analyzed using the software provided by the device manufacturer. The thermal cycling conditions were 40 cycles at 95°C for 5 s, 60°C for 30 s, and 72°C for 1 min. All RT-PCR experiments were performed in triplicate, and the specificity of each product was verified by melting curve analysis. The calculated gene expression levels were normalized to GAPDH mRNA levels.

Cytokine measurements

Cytokine concentrations were measured using human enzyme-linked immunosorbent assay (ELISA) kits for IL-6 and IL-8 (R&D Systems, Minneapolis, MN, USA) in HGF culture supernatant using the experimental procedures recommended by the manufacturer. All experiments were performed in triplicate, and data are presented as mean \pm standard deviation.

Preparation of whole-cell and nuclear extracts

HGFs (2×10^5 cells/mL) were treated with or without inactivated EBV. The cells were then washed with cold PBS and resuspended in lysis buffer (Cell Signaling Technology, Inc., Danvers, MA, USA) and protease inhibitors (Roche Diagnostics GmbH, Mannheim, Germany), incubated on ice for 10 min, and centrifuged for 15 min at 15,000 rpm. The supernatant was then collected (whole-cell extract) and stored at -80°C until use. To prepare the nuclear extract, precipitated cells were resuspended in cytoplasmic lysis buffer (Chemicon International, Temecula, CA) and incubated for 15 min on ice. The cells were vortexed and then centrifuged for 10 min at 15,000 rpm, and the supernatant was removed. The cell pellets were washed twice with cytoplasmic buffer to remove any trace of proteins from the cytoplasmic extracts, resuspended in nuclear lysis buffer (Chemicon International, Temecula, CA), and incubated on ice for 15 min. The cell suspensions were then sonicated for 10 s and centrifuged for 15 min at 15,000 rpm, and the supernatant fractions were stored at -80°C . Pierce Microplate BCA Protein Assay kit-Reducing Agent Compatible kits (Thermo Fisher Scientific, Inc.) were used to standardize the protein concentration in all samples.

Western blot analysis

The experimental procedures for western blot analysis were performed as previously described (24, 25). Briefly, equal amounts of the protein samples (20 μ g) were separated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis and transferred to a polyvinylidene fluoride membrane (EMD Millipore Corporation, Billerica, MA, USA). The membrane was probed with the appropriate antibodies, and immunoreactive proteins were visualized using SuperSignal West Pico enhanced chemiluminescence kits (Thermo Fisher Scientific, Inc.). Protein bands were detected using the ChemiDoc XRS System (Bio-Rad, Hercules, CA, USA).

Transfection and luciferase assay

HGFs (1×10^5 cells/mL) were transfected with 20 ng of pGL3- κ B luc using FuGENE 6 transfection reagent (Roche Diagnostics) according to the manufacturer's protocol (24, 25). Subsequently, 24 h after transfection, the cells were treated with EBV for an additional 8 h. These cells were then harvested using Passive Lysis Buffer (Promega Corporation, Madison, WI, USA), and the extracts were assessed for luciferase activity using the Dual-Luciferase Assay System (Promega) as previously described (24, 25). All experiments were performed in triplicate, and the data are presented as the fold increase in luciferase activity [mean \pm standard deviation (SD)] relative to controls.

Osteoclast formation and TRAP staining

Raw264.7 cells were seeded in 96-well culture plates (4×10^3 cells/well). After culturing overnight, medium was replaced with 200 μ L of α -minimal essential medium containing 10% FBS and 50 ng/mL soluble RANKL (R&D Systems) in the absence or presence of the indicated copy numbers of EBV. The cultures were then incubated at 37°C for 4 days. The

medium was replaced every other day. Cells were fixed with 4% paraformaldehyde and stained for tartrate-resistant acid phosphatase (TRAP) activity as described previously (26).

Results

Effect of EBV on IL-6 and IL-8 mRNA expression in gingival fibroblasts

To investigate the effect of EBV on the expression of IL-6 and IL-8 from HGFs, this study performed RT-PCR analysis. Because the data detected EBV DNA in deeper probing pocket depths sites (≥ 5 mm) ranging between 3.74×10^3 and 2.83×10^9 copies/mL in Japanese periodontitis patients (15), 5000 copies/mL of EBV was used to treat HGFs for different periods of time. The extent of stimulation was comparable with that elicited by TNF- α as a positive control. EBV substantially increased the expression of IL-6 and IL-8 mRNA (Fig. 1). The IL-6 mRNA level was upregulated in HGFs 4 h after incubation with inactivated EBV and reached a maximum after 12 h of stimulation. High time-dependent expression of IL-8 mRNA was observed in response to incubation with EBV.

EBV promoted the production of IL-6 and IL-8 in HGFs

The study next investigated the effect of EBV on the production of IL-6 and IL-8. HGFs were incubated with or without inactivated EBV for 48 h, and the amounts of both cytokines were measured using ELISA. As shown in Fig. 2A, the production of both IL-6 and IL-8 was induced by EBV in HGFs in a copy-dependent manner. Although CMV alone could induce the production of IL-6 and IL-8, significantly more inflammatory cytokines were produced by treatment with EBV than by treatment with CMV.

LPS, the major pathogenic component of the cell wall of *P. gingivalis*, can activate host immune responses, including the production of inflammatory cytokines and chemokines, leading to the progression of periodontal disease (1). LPS from *E. coli* and LTA from *S. aureus* were employed as stimulators of the production of inflammatory cytokines in HGFs. To compare the effects of EBV on HGFs, this work also tested LPS and LTA. As shown in Fig. 2B, the levels of IL-6 and IL-8 in HGFs were highest when the cells were

treated with EBV, rather than with *P. gingivalis* LPS, *E. coli* LPS, or LTA. These findings suggested that EBV contributes to the progression of periodontal disease by promoting the production of inflammatory cytokines.

NF- κ B was involved in the production of IL-6 and IL-8 induced by inactivated EBV in HGFs

NF- κ B is an inducible cellular transcriptional factor that regulates a wide variety of cellular genes involved in the control of the inflammatory and immune responses (27). NF- κ B is normally present in the cytoplasm in association with its inhibitory protein I κ B. Upon stimulation, signaling events are initiated, leading to the activation of the I κ B kinase (IKK) complex, which promotes the phosphorylation of I κ B on two specific serine residues (Ser32/36) in I κ B α . This phosphorylation leads to the degradation of I κ B by the 26S proteasome and thus nuclear translocation of NF- κ B (27). Because NF- κ B is part of one of the most important pathways mediating the generation of IL-6 and IL-8 (27, 28), this study examined whether inactivated EBV can activate NF- κ B, and if so, whether NF- κ B is involved in the expression of these cytokines induced by inactivated EBV. To assess whether EBV can activate NF- κ B, gingival fibroblasts were stimulated with inactivated EBV for different times and analyzed for I κ B α and NF- κ B p65 expression using western blot analysis. As shown in Fig. 3A, degradation of I κ B α was observed as early as 15 min after stimulation of these cells by inactivated EBV. These results also observed that inactivated EBV stimulated the nuclear translocation of NF- κ B p65. To further examine whether NF- κ B activation by inactivated EBV in HGFs occurred at the transcriptional level, this study employed luciferase assays using a reporter plasmid whose expression is controlled by NF- κ B. Fig. 3B shows that inactivated EBV stimulated the activation of NF- κ B-dependent genes,

such as 4xκB-luc, in a copy number-dependent manner. These results suggest that EBV could activate the NF-κB pathway in HGFs.

This study further examined whether NF-κB is involved in the induction of the expression of IL-6 and IL-8 in HGFs by inactivated EBV. Subsequently, the study used an NF-κB selective inhibitor, BAY11-70582. Gingival fibroblasts were preincubated with various concentrations of BAY11-70582 before stimulation with inactivated EBV, and the amounts of these cytokines were determined (Fig. 3C). The results found that BAY11-70582 inhibited the production of IL-6 and IL-8 induced by EBV. These results showed that NF-κB signaling contributed to the EBV-induced production of IL-6 and IL-8 in HGFs.

EBV promoted RANKL-induced osteoclast differentiation

Osteoclasts originate from the monocyte/macrophage lineage and demineralize and degrade the bone matrix (19). Osteoclast differentiation is regulated by the binding of RANK present on the surface of preosteoclasts to its ligand RANKL, which leads to the activation of NF-κB (19). To examine the effect of EBV on RANKL-induced osteoclastogenesis, macrophage-like RAW264.7 cells were treated with different copy numbers of EBV in the presence of recombinant RANKL. Although RANKL stimulation induced the formation of osteoclast-like TRAP-positive multinuclear giant cells, interestingly, EBV treatment upregulated their number in a copy number-dependent manner (Fig. 4A). Higher magnification images also revealed that EBV increased both the number and size of osteoclast-like cells in a copy number-dependent manner (Fig. 4B). In contrast, CMV were unable to stimulate formation of osteoclast-like TRAP-positive multinuclear giant cells (Fig. 4C, D)

Discussion

Although there is evidence for an association between EBV and periodontitis, the way in which EBV affects the pathogenesis and progression of periodontitis has not been well understood. It is known that excessive and/or continuous production of inflammatory cytokines occurs in response to the presence of periodontopathic bacteria and their products, such as LPS, in the periodontal tissue of periodontitis patients and is an important determinant of the onset and progression of periodontal disease. To clarify this issue, it is important to identify and characterize any complex proinflammatory cytokine network involved in periodontal pathogenesis and its link to EBV. In the present study, the effects of inactivated EBV on the expression of inflammatory cytokines in HGFs was investigated. The results demonstrated that EBV could activate NF- κ B and could subsequently promote the production of IL-6 and IL-8 in HGFs.

IL-6 activity is involved in the stimulation of acute phase protein synthesis, leukocyte recruitment, B-cell differentiation, and T-cell activation in many chronic inflammatory diseases (29). IL-8, a potent neutrophil chemoattractant and activator, has been associated with the pathogenesis of periodontal diseases and the accumulation and degranulation of neutrophils, with subsequent destruction of normal tissue (30). These cytokines could also stimulate bone resorption by promoting the formation of osteoclasts (31). Many investigators have detected IL-6 and IL-8 in gingival cavity fluid (GCF), and their levels in GCF are closely associated with the severity of inflammatory responses and periodontal tissue destruction (31–36). Reinhardt et al. reported that because IL-6 levels in GCF of periodontitis patients were higher than IL-1 levels, IL-6 plays an important role in the progression of periodontal disease (36). It has been shown that, primed by IL-1 β , IL-8 markedly upregulates neutrophils for elastase release, giving rise to a significant amplification of periodontal

inflammation (35). In the study, it was observed that the production of IL-6 and IL-8 induced by EBV was significantly stronger than that induced by *P. gingivalis* LPS. Because these inflammatory cytokines are multifunctional and exert their effects in a paracrine and autocrine fashion to modulate the inflammatory and immune responses of HGFs, these findings suggested that EBV contributes to the progression of periodontal disease by promoting the production of inflammatory cytokines.

A number of highly inducible genes encoding cytokines and chemokines contain NF- κ B-binding sites in their proximal promoters (27), and activation of NF- κ B plays a central role in virus-dependent cytokine expression and pathology. The results observed that EBV could stimulate the degradation of I κ B α and the phosphorylation of NF- κ B p65 and NF- κ B-dependent genes such as 4 \times κ B-luc. Inhibition of NF- κ B reduced the production of IL-6 and IL-8 induced by inactivated EBV in HGFs. Although the study data suggested that NF- κ B was involved in EBV-induced cytokine production in HGFs, EBV also results in the activation of other transcription factors, as well as members of signaling pathways, such as mitogen-activated protein kinases. Thus, further study is needed to understand what kind of signaling pathway is involved in the induction of inflammatory cytokines by EBV in HGFs. Because inactivated EBV was used for the stimulation of HGFs, the results suggest that infection with EBV is not required for the induction of inflammatory cytokines and NF- κ B activation in HGFs. EBV infects B cells and replicates in the epithelial cells of the oropharynx, suggesting that EBV can interact with other cell types, such as T cells and thymocytes (7, 16). EBV can also interact with monocytes and neutrophils and can modulate the expression of inflammatory cytokines such as IL-1, IL-8, macrophage inflammatory protein-1 α , and granulocyte-macrophage colony-stimulating factor without penetrating into these cells (20–23). Tanner et al. reported that EBV glycoproteins gp350 and gp220 induced the production of IL-6, so contact between the structural proteins of the virus, such as

glycoproteins, and the cells might be necessary for the induction of cytokine expression by inactivated EBV (37). These observations, including the study results, suggest that even when EBV has not infected B cells, it contributes to the onset and progression of periodontitis by simple contact with and subsequent activation of macrophages, fibroblasts, and neutrophils in the vicinity of B cells, resulting in the induction of the production and release of inflammatory cytokines. Although CD21 and Toll-like receptors are also expressed in oral cells (18, 38), it is necessary to clarify which receptors are involved in EBV-induced inflammatory cytokines in HGFs.

A major clinical symptom of periodontal disease is alveolar bone loss due to excessive resorption by osteoclasts (1, 18). *P. gingivalis* affects osteoclast formation in various ways, such as by interacting directly with osteoclast precursors, including bone marrow macrophages, or indirectly by activating HGFs and osteoblasts, which aid osteoclast formation (18). LPS is also involved in osteoclast differentiation via inducing RANKL expression (18, 19). In this study, it was found that EBV could induce the differentiation of RAW264.7 cells into osteoclast-like cells. Most recently, the data also reported that the human immune response to EBV infection may induce human osteoclast activation and cause erosive arthritis in humanized NOD/Shi-*scid*/*IL-2R γ* ^{null} (hu-NOG) mice (39). In addition, the study observed that EBER and osteoclasts were detected in the periodontium of EBV-infected hu-NOG mice (data not shown). These observations, including the results of this study, suggest that EBV may be involved in the onset and progression of inflammatory bone diseases, such as periodontal disease and rheumatoid arthritis. Although EBV does not infect mice, presenting a challenge for biomedical research, the mechanism by which EBV induces osteoclast differentiation and whether EBV actually promotes alveolar bone resorption *in vivo* is an important issue that needs to be addressed in the near future.

Earlier work previously demonstrated that *P. gingivalis* can induce EBV reactivation through epigenetic regulation (8). Significantly high levels of butyric acid have been found in the saliva of periodontitis patients, which could efficiently induce the expression of the EBV lytic switch activator *BZLF1* (40). It is assumed that the interaction between EBV and periodontopathic bacteria leads to the following negative chain of pathological events in the oral cavity: 1) poor oral hygiene causes the accumulation of butyric acid-producing periodontopathic anaerobic bacteria, such as *P. gingivalis* and *F. nucleatum*; 2) butyric acid induces EBV reactivation; 3) EBV induces the production of high concentrations of inflammatory cytokines in various types of cells and the formation of osteoclasts without infection; 4) progression of gingival inflammation and alveolar bone resorption; and 5) periodontitis escalation.

Conclusion

This study aimed to examine the effect of inactivated EBV on the expression of inflammatory cytokines in HGFs and induction of osteoclast differentiation. The conclusions of the study are as follows:

- (1) Extremely high levels of IL-6 and IL-8 were induced by inactivated EBV in a copy-dependent manner in HGFs.
- (2) NF- κ B was involved in the production of IL-6 and IL-8 induced by inactivated EBV in HGFs.
- (3) EBV promoted RANKL-induced RAW264.7 cell differentiation into osteoclast-like cells.

References

1. Holt SC and Ebersole JL: *Porphyromonas gingivalis*, *Treponema denticola*, and *Tannerella forsythia*: The “red complex,” a prototype polybacterial pathogenic consortium in periodontitis. *Periodontol 2000* 38(1): 72-122, 2005.
2. Slots J, Saygun I, Sabeti M and Kubar A: Epstein–Barr virus in oral diseases. *J Periodontal Res* 41(4): 235-244, 2006.
3. Abusleme L, Dupuy AK, Dutzan N, Silva N, Burleson JA, Strausbaugh LD, Gamonal J and Diaz PI: The subgingival microbiome in health and periodontitis and its relationship with community biomass and inflammation. *ISME J* 7(5): 1016-1025, 2013.
4. Ledder RG, Gilbert P, Huws SA, Aarons L, Ashley MP, Hull PS and McBain AJ: Molecular analysis of the subgingival microbiota in health and disease. *J Appl Environ Microbiol* 73(2): 516-523, 2007.
5. Riep B, Edesi-Neuss L, Claessen F, Skarabis H, Ehmke B, Flemmig TF, Bernimoulin JP, Göbel UB and Moter A: Are putative periodontal pathogens reliable diagnostic markers? *J Clin Microbiol* 47(6): 1705-1711, 2009.
6. Imai K and Ogata Y: How dose Epstein–Barr virus contribute to chronic periodontitis? *Int J Mol Sci* 21(6): E1940, 2020.
7. Thorley-Lawson DA and Gross A: Persistence of the Epstein–Barr virus and the origins of associated lymphomas. *N Engl J Med* 350(13): 1328-1337, 2004.
8. Imai K, Inoue H, Tamura M, Cueno ME, Inoue H, Takeichi O, Kusama K, Saito I and Ochiai K: The periodontal pathogen *Porphyromonas gingivalis* induces the Epstein–Barr virus lytic switch transactivator ZEBRA by histone modification. *Biochimie* 94(3): 839-846, 2012.
9. Contreras A, Nowzari H and Slots J: Herpesviruses in periodontal pocket and gingival tissue specimens. *Oral Microbiol Immunol* 15(1): 15-18, 2000.
10. Lu H, Zhu C, Li F, Xu W, Tao D and Feng X: Putative periodontopathic bacteria and herpesviruses in pregnant women: a case–control study. *Sci Rep* 6: 27796, 2016.
11. Slots, J. Herpesviral-bacterial interactions in periodontal diseases. *Periodontol 2000*, 52, 117-140.
12. Saygun I, Kubar A, Özdemir A and Slots J: Periodontitis lesions are a source of salivary cytomegalovirus and Epstein–Barr virus. *J Periodontal Res* 40(2): 187-191, 2005.

13. Konstantinidis A, Sakellari D, Papa A and Antoniadis A: Real-time polymerase chain reaction quantification of Epstein–Barr virus in chronic periodontitis patients. *J Periodontal Res* 40(4): 294-298, 2005.
14. Kato A, Imai K, Ochiai K and Ogata Y: Higher prevalence of Epstein–Barr virus DNA in deeper periodontal pockets of chronic periodontitis in Japanese patients. *PLoS One* 8(8): e71990, 2013.
15. Kato A, Imai K, Ochiai K and Ogata Y: Prevalence and quantitative analysis of Epstein–Barr virus DNA and *Porphyromonas gingivalis* associated with Japanese chronic periodontitis patients. *Clin Oral Investig* 19(7): 1605-1610, 2015.
16. Vincent-Bugnas S, Vitale S, Mouline CC, Khaali W, Charbit Y, Mahler P, Prêcheur I, Hofman P, Maryanski JL and Doglio A: EBV infection is common in gingival epithelial cells of the periodontium and worsens during chronic periodontitis. *PLoS One* 8(2): e80336, 2013.
17. Sunde PT, Olsen I, Enersen M and Grinde B: Patient with severe periodontitis and subgingival Epstein–Barr virus treated with antiviral therapy. *J Clin Virol* 42(2): 176-178, 2008.
18. Hajishengallis G: Periodontitis: From microbial immune subversion to systemic inflammation. *Nat Rev Immunol* 15(1): 30-44, 2015.
19. Mundy GR: Osteoporosis and inflammation. *Nutr Rev* 65(12 Pt 2): S147-S151, 2007.
20. Roberge CJ, McColl SR, Laroche B and Gosselin J: Granulocyte-macrophage colony-stimulating factor enhances EBV-induced synthesis of chemotactic factors in human neutrophils. *J Immunol* 160(5): 2442-2448, 1998.
21. Gaudreault E, Fiola S, Olivier M and Gosselin J: Epstein–Barr virus induces MCP-1 secretion by human monocytes via TLR2. *J Virol* 81(15): 8016-8024, 2007.
22. Roberge CJ, Laroche B, Rola-Pleszczynski M and Gosselin J: Epstein–Barr virus induces GM-CSF synthesis by monocytes: Effect on EBV-induced IL-1 and IL-1 receptor antagonist production in neutrophils. *Virology* 238(2): 344-352, 1997.
23. Roberge CJ, Poubelle PE, Beaulieu AD, Heitz D and Gosselin J: The IL-1 and IL-1 receptor antagonist (IL-1Ra) response of human neutrophils to EBV stimulation. Preponderance of IL-1Ra detection. *J Immunol* 156(12): 4884-4891, 1996.
24. Imai K, Togami H and Okamoto T: Involvement of histone H3 lysine 9 (H3K9) methyltransferase G9a in the maintenance of HIV-1 latency and its reactivation by BIX01294. *J Biol Chem* 285(22): 16538-16545, 2010.

25. Watanabe N, Nodomi K, Koike R, Kato A, Takeichi O, Kotani AI, Kaneko T, Sakagami H, Takei M, Ogata Y, Sato S and Imai K: EBV Imp1 in gingival epithelium potentially contributes to human chronic periodontitis via inducible IL-8 production. *In Vivo* 33(6): 1793-1800, 2019.
26. Hayata M, Watanabe N, Kamio N, Tamura M, Nodomi K, Tanaka K, Iddamalgoda A, Tsuda H, Ogata Y, Sato S, Ueda K and Imai K: Cynaropicrin from *Cynara Scolymus* L. suppresses *Porphyromonas gingivalis* LPS-induced production of inflammatory cytokines in human gingival fibroblasts and RANKL-induced osteoclast differentiation in RAW264.7 cells. *J Nat Med* 73(1): 114-123, 2019.
27. Hayden MS and Ghosh S: Signaling to NF- κ B. *Genes Develop* 18(18): 2195-2224, 2004.
28. Mukaida N: Interleukin-8: An expanding universe beyond neutrophil chemotaxis and activation. *Int J Hematol* 72(4): 391-398, 2000.
29. Hunter CA and Jones SA: IL-6 as a keystone cytokine in health and disease. *Nat Immunol* 16(5): 448-457, 2015.
30. Bickel M: The role of interleukin-8 in inflammation and mechanisms of regulation. *J Periodontol* 64(5) Supplement: 456-460, 1993.
31. Hwang YS, Lee SK, Park KK and Chung WY: Secretion of IL-6 and IL-8 from lysophosphatidic acid-stimulated oral squamous cell carcinoma promotes osteoclastogenesis and bone resorption. *Oral Oncol* 48(1): 40-48, 2012.
32. Tsai CC, Ho YP and Chen CC: Levels of interleukin-1 beta and interleukin-8 in gingival crevicular fluids in adult periodontitis. *J Periodontol* 66(10): 852-859, 1995.
33. Jin L, Söder B and Corbet EF: Interleukin-8 and granulocyte elastase in gingival crevicular fluid in relation to periodontopathogens in untreated adult periodontitis. *J Periodontol* 71(6): 929-939, 2000.
34. Bozkurt FY, Berker E, Akkuş S and Bulut S: Relationship between interleukin-6 levels in gingival crevicular fluid and periodontal status in patients with rheumatoid arthritis and adult periodontitis. *J Periodontol* 71(11): 1756-1760, 2000.
35. Brandolini L, Sergi R, Caselli G, Boraschi D, Locati M, Sozzani S and Bertini R: Interleukin-1 β primes interleukin-8-stimulated chemotaxis and elastase release in human neutrophils via its type I receptor. *Eur Cytokine Netw* 8(2): 173-178, 1997.
36. Reinhardt RA, Masada MP, Kaldahl WB, DuBois LM, Kornman KS, Choi JI, Kalkwarf KL and Allison AC: Gingival fluid IL-1 and IL-6 levels in refractory periodontitis. *J Clin Periodontol* 20(3): 225-231, 1993.

37. Tanner JE, Alfieri C, Chatila TA and Diaz-Mitoma F: Induction of interleukin-6 after stimulation of human B-cell CD21 by Epstein–Barr virus glycoproteins gp350 and gp220. *J Virol* 70(1): 570-575, 1996.
38. Kobayashi I, Shima K, Saito I, Kiyoshima T, Matsuo K, Ozeki S, Ohishi M and Sakai H: Prevalence of Epstein–Barr virus in oral squamous cell carcinoma. *J Pathol* 189(1): 34-39, 1999.
39. Nagasawa Y, Takei M, Iwata M, Nagatsuka Y, Tsuzuki H, Imai K, Imadome KI, Fujiwara S and Kitamura N: Human osteoclastogenesis in Epstein–Barr virus-induced erosive arthritis in humanized *NOD/Shi-scid/IL-2R γ null* mice. *PLoS One* 16(4): e0249340, 2021.
40. Koike R, Nodomi K, Watanabe N, Ogata Y, Takeichi O, Takei M, Kaneko T, Tonogi M, Kotani AI and Imai K: Butyric acid in saliva of chronic periodontitis patients induces transcription of the EBV lytic switch activator *BZLF1*: A Pilot study. *In Vivo* 34(2): 587-594, 2020.




Figure 1. Effects of EBV on the expression of IL-6 and IL-8 mRNA in HGFs

HGFs were treated with inactivated EBV (5000 copies/mL) or TNF- α (1 ng/mL) for the indicated times. “-” indicates treatment with virus-treated stock solution as a control. RT-PCR analysis was performed to detect the expression of IL-6 and IL-8 mRNA with specific primers. IL-6 and IL-8 mRNA levels were normalized to glyceraldehyde-3-phosphate dehydrogenase (GAPDH) mRNA levels, and they are shown as fold increase.




Figure 2. EBV-induced the release of IL-6 and IL-8 from HGFs

(A) HGFs were treated with inactivated EBV or CMV at the indicated copy numbers (copies/mL) for 48 h. Culture supernatants were collected, and IL-6 and IL-8 levels were measured using ELISA. (B) HGFs were treated with inactivated EBV (5000 copies/mL), CMV (5000 copies/mL), *P. gingivalis* LPS (500 ng or 1 µg/mL), *E. coli* LPS (100 or 200 ng/mL), or *S. aureus* LTA (5 or 10 µg/mL) for 48 h, and then, IL-6 and IL-8 levels in the supernatants were detected using ELISA. Experiments were performed in triplicate, and data are presented as mean ± standard deviation (pg/mL).




Figure 3. Activation of NF-κB by EBV and its involvement in inactivated EBV-induced IL-6 and IL-8 production

(A) Inactivated EBV-induced degradation of IκBα and translocation of p65 to the nucleus. HGFs were treated with inactivated EBV (5000 copies/mL) for the indicated time periods. Cytoplasmic extracts were fractionated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis and immunoblotted with anti-IκBα, and GAPDH antibodies. Nuclear extracts were immunoblotted with anti-p65, and PCNA antibodies. (B) Activation of NF-κB-dependent gene expression by inactivated EBV. The 4xκB-luc reporter plasmid at 20 ng was transfected into HGFs, which were incubated for 24 h and stimulated with the indicated amount of inactivated EBV for an additional 8 h. The luciferase activity of each cell lysate was then measured. All experiments were performed in triplicate, and the data are presented as the fold increase in luciferase activity (mean ± standard deviation) relative to the control. (C) Effects of BAY11-7082 on inactivated EBV-induced IL-6 and IL-8 production. HGFs were pretreated with various concentrations of BAY11-7082 for 1 h and then incubated with inactivated EBV (5000 copies/mL) for an additional 24 h. The cell culture supernatants were analyzed for IL-6 and IL-8 levels using ELISA.




Figure 4. Induction of RANKL-induced osteoclast differentiation by EBV

Raw264.7 cells were seeded in a 96-well plate and treated with recombinant RANKL (50 ng/mL) in the absence or presence of the indicated copy numbers of EBV or CMV (copies/mL) for 4 days, followed by TRAP staining. (A, C) The number of TRAP-positive multinucleated cells present in each well was counted. “VS” indicates treatment with virus-treated stock solution. (B, D) TRAP-positive multinucleated cells were observed by light microscopy.